



Anti Human Macrophage Migration Inhibitory Factor (MIF) Purified IgG Biotinylated

Product Information

Code	A00262-01-50B
Name	Anti Human MIF IgG Biotinylated Polyclonal
Clone No.	
Lot No.	
Size	50 µg
Species	Human
Host	Rabbit
Immunogen	Human MIF.
Ab Type	Purified IgG
Purification	Protein A
Formulation	Lyophilized form without preservatives free
Carry	
Storage	-20 ° C
Specificity	Human
Reconstitution	50 µl
Application	IHC WB ELISA

AVISCERA BIOSCIENCE INC.
2348 Walsh Ave. Suite C
Santa Clara, CA 95051
USA
Tel: (408) 982 0300
Fax: (408) 982 0301
Email:
Sales@AvisceraBioscience.com
www.AvisceraBioscience.com
www.AvisceraBioscience.net

Preparation

This antibody was produced from a rabbit immunized with purified recombinant human MIF. That antibody was purified by Protein A affinity and conjugated with water soluble biotin.

Formulation

50 µg of Rabbit Anti Human MIF IgG Biotinylated in 100 µl of PBS without preservatives was lyophilized.

Reconstitution

Add 50 µl of PBS to the vial to prepare antibody stock solution at 1 µg/ µl. Store reconstituted antibody at 2 to 8 ° C for up a few weeks. This antibody can also be aliquotted (by 10 µL per vial) and stored frozen at -20° C to -70° C **in a manual defrost freezer** for up 12 months without detectable loss of activity.

Storage

Lyophilized antibody can be stored at 2 ~8 ° C for a few weeks or at -20 ° C for six months. **Avoid repeated freeze-thaw cycles.**

Specificity

This antibody has been selected for its ability to recognize human MIF on indirect ELISA, Immunohistochemistry and western blot.

Applications

Indirect ELISA - This antibody can be used at 0.25 µg/ml with the appropriate secondary reagents to detect recombinant human MIF.

Western Blot - This antibody can be used at 2 µg/ml with the appropriate secondary reagents to detect recombinant human MIF under reduce conditions. The detection limit for rh MIF approximately 15 ng/lane under reducing conditions.

Immunohistochemistry - This antibody can be used at 2-4 µg/ml with the appropriate secondary antibody to detect MIF in the paraffin embedded human liver tissues (ABC).

Optimal dilutions should be determined by each laboratory for each application.